

AVISCERA BIOSCIENCE

Anti-Human Osteopontin (OPN) Monoclonal Antibody

Product Information

Code A00759-05-100

Name Anti Human OPN Mab

Clone No. 1H5G12

Lot No.

Size 100 μg

Species Human

Host Mice

Immunogen OPN (H), rec.

Ab Type IgG

Purification Protein G

Formulation lyophilized

Form without preservatives

Carry free

Storage

-20 ° C

-

Specificity Human

Reconstitu-tion $\;\;$ PBS, 100 μl

Application WB, E

AVISCERA BIOSCIENCE INC. 2348 Walsh Ave. Suite C Santa Clara, CA 95051 Tel: (408) 982 0300 Fax: (408) 982 0301

Email:

Sales@AvisceraBioscience.com www.AvisceraBioscience.com

Preparation

This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, E. coli-derived, recombinant human Osteoponin, His Tag on N-Terminal. That antibody was purified by Protein G affinity.

Formulation

Lyophilized from a 100 μg of Anti Human OPN Monoclonal Antibody in 100 μl of solution in phosphate-buffered saline (PBS) .

Reconstitution

Add 100 μ l of PBS to the vial to prepare antibody stock solution at 100 μ g /100 μ l. Store reconstituted antibody at 2 to 8 °C for up a few weeks. This antibody can also be aliquotted (by 10 uL per vial) and stored frozen at -20°C to -70°C in a manual defrost freezer for up six months without detectable loss of activity.

Storage

Lyophilized antibody can be stored at 2 $^{\circ}$ 8 $^{\circ}$ C for a few weeks or at -20 $^{\circ}$ C for 12 months. Avoid repeated freeze-thaw cycles.

Specificity

This antibody has been selected for its ability to recognize human OPN on indirect ELISA, western blot.

Applications

Indirect ELISA - This antibody can be used as antibody at 1: 8000 (0.125 μ g/ml) with the appropriate detection antibody to detect recombinant human OPN on indirect ELISA.

Western Blot-This antibody can be used as a primary antibody at 2-4 μ g/ml combine with secondary Antibody to detect human OPN on human plasma or serum under reduced condition. The positive bands were two bands, full length OPN at ~60-65KD and MMP-cleaved at ~35 – 40 KD.

Optimal dilutions should be determined by each laboratory for each application.